

SQUARE-FREE RINGS AND THEIR AUTOMORPHISM GROUP

MARTIN MONTGOMERY

ABSTRACT. Finite-dimensional square-free K -algebras have been completely characterized by Anderson and D'Ambrosia as certain semigroup algebras $A \cong K_\xi S$ over a square-free semigroup S twisted by some $\xi \in Z^2(S, K^*)$, a two-dimensional cocycle of S with coefficients in the group of units K^* of K . D'Ambrosia extended the definition of square-free to artinian rings with unity and showed every square-free ring has an associated division ring D and square-free semigroup S . We show a square-free ring R can be characterized as a semigroup ring over a square-free semigroup S twisted by some $(\alpha, \xi) \in Z^2(S, D^*)$, a two-dimensional cocycle of S with coefficients in the nonabelian group of units D^* of a division ring D . Also, to each square-free ring $R \cong D_\xi^\alpha S$ there exists a short exact sequence

$$1 \longrightarrow H_{(\alpha, \xi)}^1(S, D^*) \longrightarrow \text{Out } R \longrightarrow \text{Stab}_{(\alpha, \xi)}(\text{Aut } S) \longrightarrow 1.$$

connecting the outer automorphisms of R to certain cohomology groups related to S and D .

INTRODUCTION

The study of outer automorphisms of certain algebraic structures, beginning with Stanley [13] and continued by Scharlau [10], Baclawski [4], and Coelho [6], led to a characterization of the outer automorphism group of square-free algebras by Anderson and Dambrosia [2]. There, Clark's [5] cohomology of semigroups, was adapted to prove to every square-free algebra there is an associated short exact sequence relating the first cohomology group with the outer automorphisms of the algebra. Later, their approach was adopted by Sklar [11] to obtain results about the class of binomial algebras, which includes the class of square-free algebras.

Square-free rings were defined by D'Ambrosia [7] as a generalization of square-free algebras. She developed the basic properties of square-free rings and provided examples of square-free rings that are not of the form $D \otimes_K A$, where D is a division ring with center K and A is a square-free K -algebra.

Date: September 29, 2008.

2000 Mathematics Subject Classification. Primary: 16P20.

Key words and phrases. Square-Free Rings, Automorphism, Nonabelian Cohomology.

Presently, we will combine the above approaches to relate a square-free ring R to an associated short exact sequence with middle term the outer automorphisms of R . Utilizing ideas from non-abelian cohomology (see Dedecker [8] or Mac Lane [9]) we further extend the cohomology of semigroups from [2] and [11] to a setting related to square-free rings.

In section 1, we review some of the basic properties of a square-free ring R . These will include the existence of a canonical division ring D as well as an associated square-free semigroup S that plays an important role in the structure of R . Investigating the ring properties of R , we start to see a connection to non-abelian cohomology. In section 2, we develop cohomological ideas useful to characterize the structure of a square-free ring as related to the *thin 2-cohomology* $H^2(S, D^*)$ of [8]. By making use of the natural action of the semigroup automorphisms $\text{Aut } S$ on $H^2(S, D^*)$, we completely determine when two square-free rings are isomorphic. Finally, in section 3 we define several more ideas related to non-abelian cohomology and proceed to show for every square-free ring, there is a short exact sequence with middle term the outer automorphism group of R .

1. SQUARE-FREE RINGS

Throughout this paper, R will denote a basic indecomposable artinian ring with identity and with Jacobson radical $J = J(R)$. We will denote by $E = \{e_1, e_2, \dots, e_n\}$ a basic set of idempotents for R and note the modules

$$\overline{Re} \cong Re/Je \quad \text{and} \quad \overline{eR} \cong eR/eJ$$

with $e \in E$ are irredundant sets of the isomorphism classes of simple left and right modules of R . Since R is artinian, each finitely generated R -module M has a composition series. For $e_i \in E$, we denote by

$$c_i(M)$$

the multiplicity of the simple modules $\overline{Re_i} \cong Re_i/Je_i$ or $\overline{e_iR} \cong e_iR/e_iJ$ in a composition series for M .

A finitely generated indecomposable R -module is *square-free* if

$$c_i(M) \leq 1$$

for all $e_i \in E$. A finitely generated module is square-free if each of its indecomposable direct summands is square-free. A ring R is left square-free if ${}_R R$ is square-free; similarly, R is right square-free if R_R is square-free. We say R is *square-free* if it both left and right square-free.

By Theorem 1.4 of [7], when R is square-free, $e_i R e_i$ is a division ring for all $e_i \in E$ and for all $e_i, e_j \in E$,

$$\dim_{e_i R e_i}(e_i R e_j) \leq 1.$$

For R a connected ring, it follows there are ring isomorphisms $e_i Re_i \cong e_j Re_j$ for all $e_i, e_j \in E$. Therefore, we speak of *the* division ring (unique up to isomorphism) for a square-free ring R and denote it by D .

Associated to any artinian ring is a digraph called the *left quiver*. The left quiver of R is a directed graph Γ with vertex E and $h_{ij} = c_i(Je_j/J^2e_j)$ arrows

$$e_j \xrightarrow{\alpha_k} e_i \quad k = 1, \dots, h_{ij}, \quad 1 \leq i, j \leq n.$$

The *right quiver* of R is defined analogously. By Lemma 2.1 of [7], for R square-free, the left and right quivers are digraph duals. For convenience, we will focus on the left quiver Γ of R . Given any finite directed graph Γ , there is a natural semigroup associated to Γ , called the *path semigroup* of Γ . Associated to every square-free ring is an *algebra semigroup* called a *square-free semigroup* (see [11] and [7] for further details) S with zero element θ and a set of pairwise orthogonal idempotents $E \subseteq S$ such that

- (a) $S = \bigcup_{e_i, e_j \in E} e_i S e_j$; and
- (b) $|e_i S e_j \setminus \theta| \leq 1$ for each $e_i, e_j \in E$.

We may view S as a subset of R . To avoid any confusion in the notation, we will denote semigroup multiplication of s and t by $s \cdot t$ and ring multiplication of s and t by st . As a subset of R , S is not actually a semigroup, but is *nearly multiplicatively closed* in the following sense; for all $s, t \in S$, there is a unique $d \in D$ with

$$st = d(s \cdot t).$$

Since above d is uniquely determined by s and t we will designate it by $\xi(s, t)$ so that

$$st = \xi(s, t)s \cdot t.$$

Let R be a square-free ring. For convenience, we often will enumerate the idempotents in E as $e, f, g, \dots$. Note by definition of ξ , if $s = e \cdot s \cdot f$, it follows then that $\xi(s, f) = 1$ and $\xi(e, s) = 1$. Let $s = e \cdot s \cdot f \in R$. Since $\dim_{eRe} eRf = 1 = \dim_{fRf} fRf$, for any $fRf \in fRf$ there exists a unique $er'e \in eRe$ such that

$$er'es = s f r f$$

This correspondence defines a function $\alpha : fRf \longrightarrow eRe$. For a particular s we denote the map by α_s .

Remark 1.1. Let $s = esf$ and $t = ftg$. Since R is associative,

$$\begin{aligned} (es)(tgrg) &= (es)(\alpha_t(grg)t) \\ &= \alpha_s(\alpha_t(grg))st = \alpha_s(\alpha_t(grg))\xi(s, t)s \cdot t \end{aligned}$$

and

$$\begin{aligned} (es)(tgrg) &= (st)grg = (\xi(s, t)s \cdot t)grg \\ &= \xi(s, t)\alpha_{s \cdot t}(grg)s \cdot t. \end{aligned}$$

Since s, t and grg are arbitrary, α has the property that

$$\alpha_s \circ \alpha_t(x) = \xi(s, t)\alpha_{s \cdot t}(x)[\xi(s, t)]^{-1}.$$

Similar to previous treatments of square-free rings (see [7]), for $d \in D$, we will denote by ρ_d the inner automorphism of D given by

$$\rho_d : x \longrightarrow dxd^{-1}.$$

It follows then that α has the property

$$\alpha_s \circ \alpha_t = \rho_{\xi(s, t)} \circ \alpha_{s \cdot t}.$$

2. SEMIGROUP COHOMOLOGY

In [5], a cohomology with coefficient set in an abelian group was developed for a special class of algebra semigroups. This cohomology was generalized [2] and later extended [11] to all algebra semigroups. We now consider the case when the coefficient set is non-abelian.

Let D be a non-abelian group and let $\text{Aut}(D)$ be the automorphism group of D . Let $S^{<0>} = E$, and for each $n > 0$, define

$$S^{<n>} = \{(s_1, s_2, \dots, s_n) \in S^n \mid s_1 \cdot s_2 \cdots s_n \neq \theta\}.$$

For $n \geq 0$ and G an arbitrary (not necessarily abelian) group, let $F^n(S, G)$ denote the set of all functions from $S^{<n>}$ to G . These sets are groups under multiplication and are abelian exactly when G is abelian.

The following set we will designate as *2-cochains*:

$$F^1(S, \text{Aut}(D)) \times F^2(S, D^*)$$

We are especially interested in 2-cochains (α, ξ) which satisfy a ‘‘cocycle identity’’. For $s, t, u \in S^{<3>}$, consider all pairs $(\alpha, \xi) \in \mathcal{C}^2(S, C)$ satisfying the following properties:

$$\alpha_s(\xi(t, u)) \xi(s, t \cdot u) = \xi(s, t)\xi(s \cdot t, u) \quad (2.1)$$

and

$$\alpha_s \circ \alpha_t = \rho_{\xi(s, t)} \circ \alpha_{s \cdot t}. \quad (2.2)$$

We call such pairs *2-cocycles* and denote the set of all 2-cocycles by $Z^2(S, D^*)$. Note by Remark 1.1, to any square-free ring R there is a pair (α, ξ) satisfying Equation 2.2. Since R is associative, a straightforward check shows Equation 2.1 is also satisfied.

There is a group action $*$ of $F^0(S, \text{Aut}(D)) \ltimes F^1(S, D^*)$ on $Z^2(S, D^*)$ given by

$$(\mu, \eta) * (\alpha, \xi) = (\beta, \zeta)$$

$$\mu_e \circ \beta_s \circ \mu_f^{-1} = \rho_{\eta(s)} \circ \alpha_s, \quad \mu_e(\zeta(s, t)) = \eta(s)\alpha_s(\eta(t))\xi(s, t)\eta(s \cdot t)^{-1}$$

for $s = e \cdot s \cdot f$, $t = f \cdot t \cdot g \in S$. Altering slightly a definition from [8], we call the orbits under $*$ the *thin 2-cohomology* and denote these orbits by

$$H^2(S, D^*).$$

Definition 2.1. We say that two cocycles (α, ξ) and (β, ζ) are *cohomologous* if there is a $(\mu, \eta) \in F^0(S, \text{Aut}(D)) \ltimes F^1(S, D^*)$ such that for all $s, t \in S^*$

$$\mu_e \circ \beta_s \circ \mu_f^{-1} = \rho_{\eta(s)} \circ \alpha_s \quad (2.3)$$

and

$$\mu_e(\zeta(s, t)) = \eta(s)\alpha_s(\eta(t))\xi(s, t)\eta(s \cdot t)^{-1} \quad (2.4)$$

for $s = e \cdot s \cdot f$, $t = f \cdot t \cdot g \in S$. We will write $[\alpha, \xi] = [\beta, \zeta]$ for cohomologous cycles.

We say that a cocyle $(\alpha, \xi) \in Z^2(S, D^*)$ is *normal* if we have $\alpha_e = 1_D$ and $\xi(e, e) = 1$ for every $e \in E$. If (α, ξ) is a 2-cocyle, we may define

$$\eta(s) = \begin{cases} \xi(s, s)^{-1} & \text{if } s \in E \\ 1 & \text{otherwise} \end{cases}$$

Then $(1_D, \eta) * (\alpha, \xi)$ is a normal cocycle that is cohomologous with (α, ξ) . Hence, every cohomology class $[\alpha, \xi] \in H^2(S, D^*)$ has a normal representative.

Adopting the notation of [7], for square-free semigroups S as above, we write

$$R \cong D_\xi^\alpha S$$

for the left D -vector space on S^* with multiplication defined by

$$st = \begin{cases} \xi(s, t)s \cdot t & \text{if } s \cdot t \neq \theta \\ 0 & \text{otherwise} \end{cases}$$

for all $s, t \in S^*$, and

$$sd = \alpha_s(d)$$

for all $s \in S^*$ and $d \in D$. Extending the multiplication by linearity gives a square-free ring R with associated division ring D and semigroup S .

We can use the group $\text{Aut}(S)$ of semigroup automorphisms to define another action on $Z^2(S, D^*)$. Given $\phi \in \text{Aut}(S)$ and 2-cocycle $(\alpha, \xi) \in Z^2(S, D^*)$, we can, as in [1], [7] and [11], define a new 2-cocycle $(\alpha^\phi, \xi^\phi) \in Z^2(S, D^*)$ by

$$\alpha^\phi(s) = \alpha(\phi(s)) = \alpha_{\phi(s)} \quad \xi^\phi(s, t) = \xi(\phi(s), \phi(t))$$

for all $s, t \in S$ with $s \cdot t \neq \theta$.

If $(\mu, \eta) \in F^0(S, \text{Aut}(D)) \ltimes F^1(S, D^*)$ with

$$(\mu, \eta) * (\alpha, \xi) = (\beta, \zeta)$$

and we define $\mu^\phi(s) = \mu_{\phi(e)}$ and $\eta^\phi(s) = \eta(\phi(s))$ for $s = e \cdot s$ then

$$(\mu^\phi, \eta^\phi) * (\alpha^\phi, \xi^\phi) = (\beta^\phi, \zeta^\phi)$$

Hence, $[\alpha, \xi] = [\beta, \zeta]$ if and only if $[\alpha^\phi, \xi^\phi] = [\beta^\phi, \zeta^\phi]$ and $\text{Aut}(S)$ acts on $H^2(S, D^*)$.

We can then extend a result from [1], [7] and [11]:

Lemma 2.2. *Let S be a square-free semigroup, and let $[\alpha, \xi], [\beta, \zeta] \in H^2(S, D^*)$. If*

$$[\beta, \zeta] = [\alpha^\phi, \xi^\phi]$$

for some $\phi \in \text{Aut } S$, then as rings, we have

$$D_\zeta^\beta S \cong D_\xi^\alpha S.$$

It follows then, for each $(\alpha, \xi) \in Z^2(S, D^)$ there exists a normal $(\beta, \zeta) \in Z^2(S, D^*)$ with $D_\zeta^\beta S \cong D_\xi^\alpha S$.*

Proof. Since $[\beta, \zeta] = [\alpha^\phi, \xi^\phi]$, there is a pair $(\mu, \eta) \in F^0(S, \text{Aut}(D)) \ltimes F^1(S, D^*)$ such that

$$\mu_e \circ \beta_s \circ \mu_f^{-1} = \rho_{\eta(s)} \circ \alpha_s^\phi \tag{2.5}$$

and

$$\mu_e(\zeta(s, t)) = \eta(s) \alpha_s^\phi(\eta(t)) \xi^\phi(s, t) \eta(s \cdot t)^{-1} \tag{2.6}$$

are satisfied for $(s, t) \in S^{<2>}$ with $s = e \cdot s \cdot f$. Define a map $\gamma : D_\zeta^\beta S \rightarrow D_\xi^\alpha S$ by

$$\gamma(ds) = \mu_e(d) \eta(s) \phi(s)$$

and extend linearly. We claim γ is a ring isomorphism. Certainly, γ is one-to-one, onto, and preserves addition. Let $d_1 s, d_2 t \in D_\zeta^\beta S$ with $s = e \cdot s \cdot f, t =$

$f \cdot t \cdot g \in S$ and consider $\gamma(d_1 s d_2 t)$:

$$\begin{aligned}
\gamma(d_1 s d_2 t) &= \gamma(d_1 \beta_s(d_2) s t) = \gamma(d_1 \beta_s(d_2) \zeta(s, t) s \cdot t) \\
&= \mu_e(d_1 \beta_s(d_2) \zeta(s, t)) \eta(s \cdot t) \phi(s \cdot t) \\
&= \mu_e(d_1) [\mu_e(\beta_s(d_2))] [\mu_e(\zeta(s, t))] \eta(s \cdot t) \phi(s \cdot t) \\
&= \mu_e(d_1) [\mu_e \circ \beta_s \circ \mu_f^{-1}(\mu_f(d_2))] [\eta(s) \alpha_s^\phi(\eta(t)) \xi^\phi(s, t) \eta(s \cdot t)^{-1}] \eta(s \cdot t) \phi(s \cdot t) \\
&= \mu_e(d_1) [\eta(s) \alpha_s^\phi(\mu_f(d_2)) \eta(s)^{-1}] \eta(s) \alpha_s^\phi(\eta(t)) \xi^\phi(s, t) \phi(s \cdot t) \\
&= \mu_e(d_1) \eta(s) \alpha_{\phi(s)}(\mu_f(d_2)) \alpha_{\phi(s)}(\eta(t)) \xi(\phi(s), \phi(t)) \phi(s \cdot t) \\
&= \mu_e(d_1) \eta(s) \alpha_{\phi(s)}(\mu_f(d_2)) \alpha_{\phi(s)}(\eta(t)) \phi(s) \phi(t) \\
&= \mu_e(d_1) \eta(s) \phi(s) \mu_f(d_2) \eta(t) \phi(t) = \gamma(d_1 s) \gamma(d_2 t)
\end{aligned}$$

Hence, γ is a ring isomorphism. $\square$

For S a square-free semigroup and 2-cocycles $(\alpha, \xi), (\beta, \zeta) \in Z^2(S, D^*)$, we say a ring isomorphism $\gamma : D_\zeta^\beta S \rightarrow D_\xi^\alpha S$ is *normal* if $\gamma(E) = E$. Then as in [11], we have the following:

Lemma 2.3. *Let S be a square-free semigroup, let $[\alpha, \xi]$ and $[\beta, \zeta]$ be normal 2-cocycles in $Z^2(S, D^*)$, and let γ be a normal ring isomorphism from $D_\zeta^\beta S$ to $D_\xi^\alpha S$. Then there exists an automorphism ϕ of the semigroup S and $(\mu, \eta) \in F^0(S, \text{Aut}(D)) \ltimes F^1(S, D^*)$ such that*

$$\gamma(ds) = \mu_e(d) \eta(s) \phi(s)$$

for $s = e \cdot s \cdot f \in S^*$.

Proof. First, we show for every $s \in S^*$, there is a unique element $\phi(s) \in S^*$ such that $\gamma(s) \in D\phi(s)$. Since $\gamma : D_\zeta^\beta S \rightarrow D_\xi^\alpha S$ is a ring isomorphism, we must have

$$\gamma(s) = \sum_{t \in S^*} d_t t$$

with $d_t \in D$. But for some $e, f \in E$, $s = e \cdot s \cdot f$ and

$$\gamma(s) = \gamma(e) \gamma(s) \gamma(f) = \gamma(e) \left(\sum_{t \in S^*} d_t t \right) \gamma(f).$$

Since $d_t t = t \alpha_t^{-1}(d_t)$ for all d_t and t , it follows that

$$\gamma(s) = \left(\sum_{t \in S^*} d_t \gamma(e) \cdot t \cdot \gamma(f) \right).$$

But since S is square-free and γ is an isomorphism, $|\gamma(e) \cdot S^* \cdot \gamma(f)| = 1$. Denote by t_s the unique element in S^* such that $\gamma(e) \cdot t_s \cdot \gamma(f)$ whenever $s = e \cdot s \cdot f$ and define $\phi(s)$ by

$$\phi(s) = t_s$$

and $\phi(\theta) = \theta$. Since γ is a ring isomorphism, it follows that $\phi \in \text{Aut } S$. Also, $\gamma(s) = d_s \phi(s)$ for some unique $d_s \in D$. Define $\eta : S^* \rightarrow D^*$ by

$$\eta(s) = d_s.$$

Then $\eta \in F^1(S, D^*)$. Now for any $d \in D$, we must have

$$\gamma(ds) = \gamma(de \cdot s) = \gamma(de) \gamma(s) = \gamma(de) \eta(s) \phi(s) \in D \phi(s)$$

and it follows that $\gamma(de) \in D$. Define μ by

$$\mu_e(d) = \gamma(de)$$

for every $e \in E$ and d in D . Again, since γ is an isomorphism, it follows that $\mu_e \in \text{Aut}(D)$. Then $\mu \in F^0(S, \text{Aut}(D))$ and

$$\gamma(ds) = \mu_e(d) \eta(s) \phi(s)$$

as desired. $\square$

Suppose $e \in E$. As above, $\gamma(e) = \gamma(e^2) = \gamma(e) \gamma(e)$ and the map $\eta \in F^1(S, D^*)$ from Lemma 2.3 has the property

$$\eta(e) \phi(e) = \eta(e) \phi(e) = \eta(e) \phi(e) \Rightarrow \eta(e) = \eta(e) \alpha_e(\eta(e)) = [\eta(e)]^2.$$

Hence, $\eta(e) = 1$ for all $e \in E$.

We now have the following theorem:

Theorem 2.4. *Let S be a square-free semigroup, and let $[\alpha, \xi], [\beta, \zeta] \in H^2(S, D^*)$. Then*

$$D_\zeta^\beta S \cong D_\xi^\alpha S$$

if and only if there exists an automorphism $\phi \in \text{Aut}(S)$ such that

$$[\alpha^\phi, \xi^\phi] = [\beta, \zeta]$$

Proof. In Lemma 2.2, we showed that $[\alpha^\phi, \xi^\phi] = [\beta, \zeta]$ for some $\phi \in \text{Aut}(S)$ gives $D_\xi^\alpha S \cong D_\zeta^\beta S$ as rings.

Suppose $D_\xi^\alpha S \cong D_\zeta^\beta S$ via the ring isomorphism γ , and assume, without loss of generality, that ξ and ζ are normal. Since γ is an isomorphism, we know that E and $\gamma(E)$ are both basic sets for $D_\zeta^\beta S$; therefore, there exists an invertible element $u \in D_\zeta^\beta S$ such that

$$\rho_u \circ \gamma(E) = u \gamma(E) u^{-1} = E$$

(see [3], Exercise 7.2 (2)). Then $\rho_u \circ \gamma$ is a normal isomorphism from $D_\xi^\alpha S$ to $D_\zeta^\beta S$; thus we may assume without loss of generality that we begin with γ a normal map. By Lemma 2.3, there exists $(\mu, \eta) \in F^0(S, \text{Aut}(D)) \ltimes F^1(S, D^*)$ and an automorphism $\phi \in \text{Aut}(S)$ such that

$$\gamma(ds) = \mu_e(d)\eta(s)\phi(s)$$

for every $s = e \cdot s \cdot f \in S^*$. Then for $s, t \in S$ with $s \cdot t \neq \theta$ (and $\phi(s) \cdot \phi(t) \neq \theta$), we have

$$\begin{aligned} \mu_e(\zeta(s, t))\eta(s \cdot t)\phi(s \cdot t) &= \gamma(\zeta(s, t)s \cdot t) \\ &= \gamma(st) = \gamma(s)\gamma(t) \\ &= \eta(s)\phi(s)\eta(t)\phi(t) \\ &= \eta(s)\alpha_s^\phi(\eta(t))\xi^\phi(s, t)\phi(s \cdot t) \end{aligned}$$

Equating coefficients in D gives

$$\mu_e(\zeta(s, t)) = \eta(s)\alpha_s^\phi(\eta(t))\xi^\phi(s, t)\eta(s \cdot t)^{-1}. \quad (2.7)$$

For $s = e \cdot s \cdot f \in S$ and $t = f \cdot t \in S$ such that $s \cdot t \neq \theta$, consider the arbitrary element of D denoted by $\mu_f^{-1}(d)$. Then

$$\begin{aligned} \gamma(s(\mu_f^{-1}(d)f)) &= \gamma(\beta_s(\mu_f^{-1}(d))s) \Rightarrow \gamma(s)\gamma(\mu_f^{-1}(d)f) = \mu_e(\beta_s(\mu_f^{-1}(d)))\eta(s)\phi(s) \\ &\Rightarrow \eta(s)\phi(s)d = \mu_e(\beta_s(\mu_f^{-1}(d)))\eta(s)\phi(s) \\ &\Rightarrow \eta(s)\alpha_s^\phi(d) = \mu_e(\beta_s(\mu_f^{-1}(d)))\eta(s) \end{aligned}$$

It follows then that

$$\mu_e \circ \beta_s \circ \mu_f^{-1} = \rho_{\eta(s)} \circ \alpha_s^\phi \quad (2.8)$$

By Equations 2.7 and 2.8, we have

$$(\mu, \eta) * (\alpha^\phi, \xi^\phi) = (\beta, \zeta).$$

Hence $[\alpha^\phi, \xi^\phi] = [\beta, \zeta] \in H^2(S, D^*)$, as desired. $\square$

3. THE AUTOMORPHISM GROUP OF A SQUARE-FREE RING

Let R be a binomial ring. By Lemma 2.2 and Theorem 2.4, we may assume $R \cong D_\xi^\alpha S$, where S is a square-free semigroup with idempotent set E , D is the canonical division ring related to R and $(\alpha, \xi) \in Z^2(S, D^*)$ is a normal 2-cocycle. We will now consider the outer automorphism group

$$\text{Out} R = \text{Aut } R / \text{Inn } R$$

of R , where $\text{Aut } R$ is the group of all ring automorphisms and $\text{Inn}(R)$ is the normal subgroup of $\text{Aut } R$ consisting of maps ρ_r for invertible $r \in R$. Finally, 1_D and 1_S , respectively, will each denote the identity automorphism of the ring D and the semigroup S . An unlabeled 1 will denote the multiplicative identity of D .

Definition 3.1. Let $(\alpha, \xi) \in Z^2(S, D^*)$. A pair $(\mu, \eta) \in F^0(S, \text{Aut}(D)) \ltimes F^1(S, D^*)$ is called an (α, ξ) 1-cocyle if

$$\mu_e \circ \alpha_s \circ \mu_f = \rho_{\eta(s)} \circ \alpha_s \quad (3.1)$$

and

$$\mu_e(\xi(s, t)) = \eta(s)\alpha_s(\eta(t))\xi(s, t)\eta(s \cdot t)^{-1} \quad (3.2)$$

for all $s = e \cdot s \cdot f \in S$ and $(s, t) \in S^{<2>}$. The set of all (α, ξ) 1-cocyles will be denoted by $Z_{(\alpha, \xi)}^1(S, D^*)$.

Remark 3.2. Note that $Z_{(\alpha, \xi)}^1(S, D^*)$ is the stabilizer of (α, ξ) under the action $*$ of $F^0(S, \text{Aut}(D)) \ltimes F^1(S, D^*)$ on $Z^2(S, D^*)$. Hence $Z_{(\alpha, \xi)}^1(S, D^*)$ is a subgroup of $F^0(S, \text{Aut}(D)) \ltimes F^1(S, D^*)$. It easily follows that $Z_{(\beta, \zeta)}^1(S, D^*) \cong Z_{(\alpha^\phi, \xi^\phi)}^1(S, D^*)$ whenever $[\beta, \zeta] = [\alpha^\phi, \xi^\phi]$ in $H^2(S, D^*)$.

There is a group action $\star : F^0(S, D^*) \times Z_{(\alpha, \xi)}^1(S, D^*) \rightarrow Z_{(\alpha, \xi)}^1(S, D^*)$ given by

$$\epsilon \star (\mu, \eta) = (\hat{\mu}, \hat{\eta})$$

$$(\hat{\mu}(s))(d) = (\rho_{\epsilon(e)} \circ \mu_s)(d) \quad \hat{\eta}(s) = \epsilon(e)\eta(s)\alpha_s(\epsilon_f^{-1})$$

for $d \in D$ and $s = e \cdot s \cdot f$.

Definition 3.3. We call the orbit of $(1_D, 1)$ under this action the set of (α, ξ) 1-coboundaries and denote it by $B_{(\alpha, \xi)}^1(S, D^*)$. An easy check shows $B_{(\alpha, \xi)}^1(S, D^*)$ is a normal subgroup of $Z_{(\alpha, \xi)}^1(S, D^*)$. We define the (α, ξ) 1-cohomology group of S with coefficients in D^* to be the factor group

$$H_{(\alpha, \xi)}^1(S, D^*) = Z_{(\alpha, \xi)}^1(S, D^*) / B_{(\alpha, \xi)}^1(S, D^*).$$

For $(\mu, \eta) \in Z_{(\alpha, \xi)}^1(S, D^*)$, let $\sigma_{\mu\eta} : R \rightarrow R$ be the map defined by

$$\sigma_{\mu\eta}(ds) = \mu_e(d)\eta(s)s$$

for $d \in D$ and $s = e \cdot s \in S$. By extending $\sigma_{\mu\eta}$ linearly and utilizing the defining properties of $Z_{(\alpha, \xi)}^1(S, D^*)$, one can verify that $\sigma_{\mu\eta}$ is a ring homomorphism. Then we have the following:

Lemma 3.4. *There is a homomorphism $\hat{\Lambda} : Z_{(\alpha, \xi)}^1(S, D^*) \rightarrow \text{Aut } R$.*

Proof. For $(\mu, \eta) \in Z_{(\alpha, \xi)}^1(S, D^*)$ and $ds \in R \cong D_\xi^\alpha S$, we easily see

$$\begin{aligned} \hat{\Lambda}((\mu, \eta)) \circ \Lambda((\hat{\mu}, \hat{\eta}))(ds) &= \hat{\Lambda}((\mu, \eta))(\hat{\mu}(d)\hat{\eta}(s)(s)) \\ &= \mu(\hat{\mu}(d)\hat{\eta}(s))\eta(s)s \\ &= (\mu \circ \hat{\mu})(d)\mu(\hat{\eta}(s))\eta(s)s = \hat{\Lambda}((\mu, \eta)(\hat{\mu}, \hat{\eta}))(ds) \end{aligned}$$

□

Lemma 3.5. *There exists a monomorphism*

$$\Lambda : H_{(\alpha, \xi)}^1(S, D^*) \rightarrow \text{Out } R$$

defined by

$$\Lambda : (B^1(S, D^*))(\mu, \eta) \rightarrow (\text{Inn } R)\sigma_{\mu\eta}$$

for every $(\mu, \eta) \in Z_{(\alpha, \xi)}^1(S, D^)$.*

Proof. It is enough to show for $(\mu, \eta) \in Z_{(\alpha, \xi)}^1(S, D^*)$ that $(\mu, \eta) \in B_{(\alpha, \xi)}^1(S, D^*)$ if and only if $\sigma_{\mu\eta} \in \text{Inn } R$. First suppose $(\mu, \eta) \in B_{(\alpha, \xi)}^1(S, D^*)$. Then there exists $\epsilon \in F^0(S, D^*)$ such that

$$\mu_e(d) = \rho_{\epsilon(e)}(d)$$

$$\eta(s) = \epsilon(e)\alpha_s(\epsilon(f)^{-1})$$

for $d \in D$ and $s = e \cdot s \cdot f \in S$. Let $r = \sum_{e \in E} \epsilon(e)e \in R$. Then $r^{-1} = \sum_{f \in E} \epsilon(f)^{-1}f \in R$ and

$$\sigma_{\mu\eta}(ds) = \mu_e(d)\eta(s)s = \rho_{\epsilon(e)}(d)\epsilon(e)\eta(s)\alpha_s(\epsilon(f)^{-1})s$$

$$\mu_e(d)\eta(s)s = \rho_{\epsilon(e)}(d)\epsilon(e)\eta(s)s\epsilon(f)^{-1}$$

$$= \epsilon(e)ds\epsilon_f^{-1}$$

$$= \left(\sum_{e \in E} \epsilon(e)e \right) ds \left(\sum_{f \in E} \epsilon^{-1}(f)f \right)$$

$$= \rho_r(ds)$$

and it follows that $\sigma_{\mu\eta} \in \text{Inn } R$.

Now suppose $\sigma_{\mu\eta} \in \text{Inn } R$. Then for all $d \in D^*$ and $s = e \cdot s \cdot f \in S$, there exists $r \in R$ such that

$$\sigma_{\mu\eta}(ds) = \rho_r(ds) \Rightarrow \mu(d)\eta(s)s = r(ds)r^{-1}.$$

Since $r \in R$, we have $r = \sum_{t \in S} d_t t$ and $r^{-1} = \sum_{t \in S} \hat{d}_t t$. But

$$\mu(d)\eta(s)s = r(ds)r^{-1}$$

implies that

$$\eta(s)s = r(s)r^{-1} = \left(\sum_{t \in S} d_t t \right) s \left(\sum_{s \in S} \hat{d}_t t \right)$$

$$= d_e s \hat{d}_f = d_e \alpha_s(\hat{d}_f)s.$$

Since $\sigma_{\mu\eta} \in \text{Aut } R$ and $\sigma_{\mu\eta}(e) = \sigma_{\mu\eta}(e^2)$ implies

$$\eta(e) = \eta(e)\alpha_e(\eta(e)) = \eta(e)\eta(e) \Rightarrow \eta(e) = 1,$$

setting $s = e$ from above gives

$$1 = d_e \hat{d}_e \Rightarrow \hat{d}_e = d_e^{-1}$$

for all $e \in E$. Define $\epsilon \in F^0(S, D^*)$ by

$$\epsilon(e) = d_e$$

for $e \in E$. By the work above,

$$\eta(s) = \epsilon(e)\alpha_s(\epsilon(f)^{-1}) \tag{3.3}$$

and

$$\mu(d)\eta(s)s = r(ds)r^{-1} \Rightarrow \mu(d)d_e\alpha_s(d_f^{-1})s = d_ed\alpha_s(d_f^{-1})s$$

from which we have

$$\mu(d) = d_ed d_e^{-1} = \rho_{e(e)}(d). \quad (3.4)$$

Equations 3.3 and 3.4 show that $(\mu, \eta) \in B^1(S, D^*)$. $\square$

For the idempotent set $E \subset S$, define $\text{Aut}_0 R$ by

$$\text{Aut}_0 R = \{\gamma_0 \mid \gamma_0(E) = E\}.$$

For any $\gamma \in \text{Aut}(R)$, it follows (by Lemma 2.4 of [2]) there exists a normal automorphism γ_0 of R with $(\text{Inn } R)\gamma = (\text{Inn } R)\gamma_0$. If in addition, $\rho_r \in \text{Inn } R \cap \text{Aut}_0 R$, we must have $\rho_r(E) = E$. If R is basic, since $Re \cong R\rho_r(e)$ for all $e \in E$, we must have $\rho_r(e) = e$ for all $e \in E$. It follows if $(\text{Inn } R)\gamma_0 = (\text{Inn } R)\gamma'_0$ with $\gamma_0, \gamma'_0 \in \text{Aut}_0 R$ then $\gamma_0(e) = \gamma'_0(e)$ for all $e \in E$. By Lemma 2.3, we have

$$\gamma_0(e) = \phi(e) = \gamma'_0(e)$$

for some $\phi \in \text{Aut } E$. Since S is square-free, ϕ is also an automorphism of S and we have a well-defined map

$$\Phi : \text{Out } R \rightarrow \text{Aut } S$$

given by

$$(\text{Inn } R)\gamma_0 \mapsto \phi,$$

where $\gamma_0 \in \text{Aut}_0 R$ such that $\gamma_0(ds) = \mu_e(d)\eta(s)\phi(s)$ for $d \in D$ and $s = e \cdot s \in S$. A straightforward check shows that Φ is a group homomorphism.

Lemma 3.6. *Im $\Lambda = \text{Ker } \Phi$*

Proof. Let $(B^1(S, D^*))(\mu, \eta) \in H^1_{(\alpha, \xi)}(S, D^*)$. Then

$$\Phi\Lambda((B^1(S, D^*))(\mu, \eta)) = \Phi((\text{Inn}(R))\sigma_{\mu\eta}) = 1_S.$$

It follows then that $\text{Im } \Lambda \subseteq \text{Ker } \Phi$.

Now let $\gamma \in \text{Aut}_0 R$ with $(\text{Inn } R)\gamma \in \text{Ker } \Phi$. Then by Lemma 2.3, for each $s \in S$;

$$\gamma(s) = \eta(s)s.$$

Because γ is a homomorphism, $\gamma(st) = \gamma(s)\gamma(t)$ and it follows that

$$\mu_e(\xi(s, t))\eta(s \cdot t) = \eta(s)\alpha_s(\eta(t))\xi(s, t). \quad (3.5)$$

Also, $\gamma(s(dt)) = \gamma(s)\gamma(dt)$ and from Equation 3.5 we have

$$\mu_e \circ \alpha_s \circ \mu_f^{-1} = \rho_{\eta(s)} \circ \alpha_s. \quad (3.6)$$

From the above equations we see (μ, η) is a 1-cocycle and we have

$$(\text{Inn } R)\gamma = (\text{Inn } R)\sigma_{\mu\eta} = \Lambda((B^1(S, D^*))(\mu, \eta)) \in \text{Im } \Lambda.$$

$\square$

We now focus on finding the image of Λ in $\text{Aut } S$. Recall that $\text{Aut } S$ acts on $H^2(S, D^*)$. For any element $[\alpha, \xi] \in H^2(S, D^*)$, the stabilizer of the element is the set

$$\text{Stab}_{[\alpha, \xi]}(\text{Aut } S) = \{\phi \in \text{Aut}(S) \mid [\alpha, \xi] = [\alpha^\phi, \xi^\phi]\}.$$

This means there exists an element $(\mu, \eta) \in F^0(S, \text{Aut}(D)) \ltimes F^1(S, D^*)$ such that

$$\mu_e \circ \alpha_s \circ \mu_f^{-1} = \rho_{\eta(s)} \circ \alpha_s^\phi \quad (3.7)$$

and

$$\mu_e(\xi(s, t)) = \eta(s)\alpha_s(\eta(t))\xi^\phi(s, t)\eta(s \cdot t)^{-1} \quad (3.8)$$

for all $(s, t) \in S^{<2>}$.

Lemma 3.7. *Im $\Phi = \text{Stab}_{[\alpha, \xi]}(\text{Aut } S)$.*

Proof. First, suppose $\phi \in \text{Aut } S$ and $(\mu, \eta) \in F^0(S, \text{Aut}(D)) \ltimes F^1(S, D^*)$ with

$$\mu_e \circ \alpha_s \circ \mu_f^{-1} = \rho_{\eta(s)} \circ \alpha_s^\phi \quad (3.9)$$

and

$$\mu_e(\xi(s, t)) = \eta(s)\alpha_s(\eta(t))\xi^\phi(s, t)\eta(s \cdot t)^{-1} \quad (3.10)$$

for all $(s, t) \in S^{<2>}$. Define a map $\sigma_{\mu\eta\phi} : R \rightarrow R$ by

$$\sigma_{\mu\eta\phi}(ds) = \mu_e(d)\eta(s)\phi(s)$$

for $d \in S$ and $s = e \cdot s \in S$. Clearly, $\sigma_{\mu\eta\phi}$ preserves addition. We must show that it also preserves multiplication. Let $d_1, d_2 \in D$ and $s = e \cdot s \cdot f$, $t = f \cdot t \in S$ with $s \cdot t \neq \theta$. So

$$\begin{aligned} \sigma_{\mu\eta\phi}(d_1 s) \sigma_{\mu\eta\phi}(d_2 t) &= \mu_e(d_1)\eta(s)\phi(s)\mu_f(d_2)\eta(t)\phi(t) \\ &= \mu_e(d_1)\eta(s)\alpha_s^\phi(\mu_f(d_2))\alpha_s^\phi(\eta(t))\phi(s)\phi(t) \\ &= \mu_e(d_1)\rho_{\eta(s)}(\alpha_s^\phi(\mu_f(d_2)))\eta(s)\alpha_s^\phi(\eta(t))\xi^\phi(s, t)\phi(s \cdot t) \\ &= \mu_e(d_1)[\rho_{\eta(s)}(\alpha_s^\phi(\mu_f(d_2)))]\eta(s)\alpha_s^\phi(\eta(t))\xi^\phi(s, t)\eta(s \cdot t)^{-1}\eta(s \cdot t)\phi(s \cdot t) \\ &= \mu_e(d_1)[\mu_e(\alpha_s(d_2))]\mu_e(\xi(s, t))\eta(s \cdot t)\phi(s \cdot t) \\ &= \sigma_{\mu\eta\phi}(d_1 \alpha_s(d_2) \xi(s, t) s \cdot t) = \sigma_{\mu\eta\phi}(d_1 s d_2 t) \end{aligned}$$

and it follows that $\sigma_{\mu\eta\phi} \in \text{Aut}_0 R$.

Conversely, now suppose that $\gamma \in \text{Aut}_0 R$ with $\phi = \Phi(\text{Inn}(R))\gamma$. Then by Lemma 2.3, $\gamma = \sigma_{\mu\eta\phi}$ for some $\mu \in F^0(S, \text{Aut}(D))$ and $\eta \in F^1(S, D^*)$. Since

$\gamma = \sigma_{\mu\eta\phi}$ is a homomorphism, $\sigma_{\mu\eta\phi}(st) = \sigma_{\mu\eta\phi}(s)\sigma_{\mu\eta\phi}(t)$ for all $s = e \cdot s, t = f \cdot t \in S$ and

$$\sigma_{\mu\eta\phi}(st) = \sigma_{\mu\eta\phi}(s)\sigma_{\mu\eta\phi}(t) \Rightarrow \sigma_{\mu\eta\phi}(\xi(s, t)s \cdot t) = \eta(s)\phi(s)\eta(t)\phi(t)$$

$$\Rightarrow \mu_e(\xi(s, t))\eta(s \cdot t)\phi(s \cdot t) = \eta(s)\alpha_s^\phi(\eta(t))\xi^\phi(s, t)\phi(s \cdot t)$$

Equating coefficients of $\phi(s \cdot t)$, it follows

$$\mu_e(\xi(s, t)) = \eta(s)\alpha_s^\phi(\eta(t))\xi^\phi(s, t)\eta(s \cdot t)^{-1} \quad (3.11)$$

Furthermore,

$$\sigma_{\mu\eta\phi}(s\mu_f^{-1}(d)t) = \sigma_{\mu\eta\phi}(s)\sigma_{\mu\eta\phi}(\mu_f^{-1}(d)t) \Rightarrow \sigma_{\mu\eta\phi}(\alpha_s(\mu_f^{-1}(d)\xi(s, t)s \cdot t) = \eta(s)\phi(s)d\eta(t)\phi(t)$$

$$\Rightarrow \mu_e(\alpha_s(\mu_f^{-1}(d)))[\mu_e(\xi(s, t))\eta(s \cdot t)]\phi(s \cdot t) = \eta(s)\alpha_s^\phi(d)\alpha_s^\phi(\eta(t))\xi^\phi(s, t)\phi(s \cdot t)$$

Equating coefficients and using Equation 3.11 above, it follows

$$\mu_e(\alpha_s(\mu_f^{-1}(d))[\eta(s)\alpha_s^\phi(\eta(t))\xi^\phi(s, t)] = \eta(s)\alpha_s^\phi(d)\alpha_s^\phi(\eta(t))\xi^\phi(s, t)$$

$$\Rightarrow \mu_e(\alpha_s(\mu_f^{-1}(d))) = \eta(s)\alpha_s^\phi(d)\eta(s)^{-1}$$

$$\Rightarrow \mu_e \circ \alpha_s \circ \mu_f^{-1} = \rho_{\eta(s)} \circ \alpha_s^\phi.$$

So $\phi \in \text{Stab}_{[\alpha, \xi]}(\text{Aut}(S))$ and $\text{Im } \Phi = \text{Stab}_{[\alpha, \xi]}(\text{Aut}(S))$. □

Finally, we have the main theorem of this section.

Theorem 3.8. *Let R be a square-free ring with associated semigroup S , division ring D and normal $(\xi, \alpha) \in Z^2(S, D^*)$ such that $R \cong D_\xi^\alpha S$. The following sequence is exact:*

$$1 \longrightarrow H_{(\alpha, \xi)}^1(S, D^*) \xrightarrow{\Lambda} \text{Out } R \xrightarrow{\Phi} \text{Stab}_{(\alpha, \xi)}(\text{Aut } S) \longrightarrow 1.$$

If $[\alpha, \xi] = [1_D, 1]$, then $\text{Stab}_{(\alpha, \xi)}(\text{Aut } S) = \text{Aut } S$ and the following sequence is exact:

$$1 \longrightarrow H_{(\alpha, \xi)}^1(S, D^*) \xrightarrow{\Lambda} \text{Out } R \xrightarrow{\Phi} \text{Aut } S \longrightarrow 1.$$

Moreover, this last sequence splits, and so in this case $\text{Out } R$ is isomorphic to

$$H_{(\alpha, \xi)}^1(S, D^*) \ltimes \text{Aut } S.$$

Proof. By Lemma 2.2, we may assume (α, ξ) is normal. The previous lemmas of this section prove the statement regarding the first short exact sequence, so it enough to show the sequence splits in the case that $[\alpha, \xi] = [1_D, 1]$. We can

assume that $(\alpha, \xi) = (1_D, 1)$. In this case $s(dt) = ds \cdot t \in R$ for all $d \in D$ and $s, t \in S$. Given $\phi \in \text{Aut } S$, define the map σ_ϕ by

$$\sigma_\phi(ds) = d\phi(s)$$

for all $d \in D$ and $s \in S$ and extend linearly. An easy check shows that $\sigma_\phi \in \text{Aut}_0 R$. Now define a map $\Psi : \text{Aut } S \rightarrow \text{Out } R$ by

$$\Psi : \phi \rightarrow (\text{Inn } R)\sigma_\phi$$

for all $\phi \in \text{Aut } S$. A straightforward check shows Ψ to be a homomorphism of groups with $\Phi\Psi = 1_S \in \text{Aut } S$. Then the sequence is split and

$$\text{Out } R \cong H_{(\alpha, \xi)}^1(S, D^*) \rtimes \text{Aut } S.$$

□

As in the algebra case, we have the following immediate corollaries to Theorem 3.8.

Corollary 3.9. *Let $R \cong D_\xi^\alpha S$ be a square-free ring. If $H_{(\alpha, \xi)}^1(S, D^*) = 1$, then*

$$\text{Out } R \cong \text{Stab}_{[\alpha, \xi]}(\text{Aut } S).$$


Moreover, if $[\alpha, \xi] = [1_D, 1]$, then

$$\text{Out } R \cong \text{Aut } S.$$

Corollary 3.10. *Let $R \cong D_\xi^\alpha S$ be a square-free ring. Then every automorphism of R is inner if and only if*

$$H_{(\alpha, \xi)}^1(S, D^*) = \text{Stab}_{[\alpha, \xi]}(\text{Aut } S) = 1.$$

Example 3.11. Let D be a division ring that has an automorphism γ that is not inner and let S be a square-free semigroup with the following Hasse diagram



and multiplication defined by $s \cdot t \neq \theta$ only if s or t in E . Define $\alpha \in F^1(S, \text{Aut}(D))$ by

$$\alpha_s = \begin{cases} \gamma & \text{if } s = s_{34} \\ 1 & \text{otherwise} \end{cases}$$

and take $\xi = 1$. For $s = s_{ij}$ denote the maps α_s and μ_s by α_{ij} and μ_{ij} respectively.

By Corollary 3.5 of [7], $R = D_\xi^\alpha S$ is a square-free ring not isomorphic to $D \otimes_K A$, where K is the center of D and A is a square-free K -algebra.

Both $Z_{(\alpha, \xi)}^1(S, D^*)$ and $B_{(\alpha, \xi)}^1(S, D^*)$ can be determined using the semi-group structure of S and the defining properties of each group. For $(\mu, \eta) \in Z_{(\alpha, \xi)}^1(S, D^*)$ we must have $\eta(e_i) = 1$ for $i = 1, 2, 3, 4$ and no restrictions on $\eta(s)$ for $s \notin E$. For μ , the maps are determined by the idempotent e for any $s = e \cdot s \cdot f \in S$. But since $\alpha_s = 1$ for all but one $s \in S$, the identity $\mu_j = \rho_{\mu_i(\eta(s_{ij}))}$ implies that μ is determined by μ_1 and our previous choices for $\eta(s)$. Then

$$Z_{(\alpha, \xi)}^1(S, D^*) = \text{Aut } D \ltimes \Pi_{i=1}^4 D^*.$$

A similar procedure for $(\mu, \eta) \in B_{(\alpha, \xi)}^1(S, D^*)$ shows that η is determined by $\epsilon : E \rightarrow D^*$ and this determines μ as well since $\mu_{ij} = \rho_{\epsilon(e_i)}$. Therefore,

$$B_{(\alpha, \xi)}^1(S, D^*) = \Pi_{i=1}^4 D^*.$$

and

$$H_{(\alpha, \xi)}^1(S, D^*) = \text{Aut } D.$$

A tedious but straightforward check shows that $\text{Stab}_{[\alpha, \xi]}(\text{Aut } S) = \text{Aut } S \cong \mathbb{Z}_2$. Hence, to R we may associate the following short exact sequence

$$1 \longrightarrow \text{Aut } D \longrightarrow \text{Out } R \longrightarrow \mathbb{Z}_2 \longrightarrow 1.$$

ACKNOWLEDGEMENT

The author would like to thank Professor Frank Anderson for his continued guidance and inspiration.

REFERENCES

- [1] F. W. ANDERSON, Transitive square-free algebras, *J. Algebra* **62** (1980), 61-85.
- [2] F.W. ANDERSON AND B.K. D'AMBROSIA, Square-free algebras and their automorphism groups, *Comm. Alg.*, **24** (1996) (10), 3163-3191.
- [3] F.W. ANDERSON AND K.R. FULLER, *Rings and Categories of Modules* (2d ed.), Springer-Verlag, New York-Berlin-Heidelberg, 1992.
- [4] K. BACLAWSKI, Automorphisms and derivations of incidence algebras, *Proc. Amer. Math. Soc.*, **36** (1972), 351-356.
- [5] W.E. CLARK, Twisted matrix units semigroup algebras, *Duke Math J.*, Sept. (1967) 417-423.
- [6] S. P. COELHO, The automorphism group of a structural matrix algebra, *Linear Algebra Appl.*, **195** (1993), 35-58.
- [7] B.K. D'AMBROSIA, Square-free rings, *Comm. Alg.*, 27(5) (1999), 2045-2071.

- [8] P. DEDECKER, Three dimensional non-abelian cohomology for groups, *1969 Category Theory, Homology Theory and their Applications, II (Battelle Institute Conference, Seattle, Wash., Vol. II)* 32-64 Springer, Berlin 1968.
- [9] S. MAC LANE, *Homology*, Springer-Verlag, New York-Berlin-Heidelberg, 1995.
- [10] W. SCHARLAU, Automorphisms and involutions of incidence algebras, in "Representations of Algebras, Ottawa, 1974", *Lecture Notes in Math. No. 488*, Springer-Verlag, New York-Berlin-Heidelberg, 1975.
- [11] J. K. SKLAR, Binomial algebras, *Comm. Alg.* , **30**(4) (2002), 1961-1978.
- [12] J. K. SKLAR, Binomial Rings, *Comm. Alg.* , **32**(4) (2004), 1385-1399.
- [13] R.P. STANLEY, Structure of incidence algebras and their automorphism groups, *Bull. Amer. Math. Soc.*, **76** (1970), 1236-1239.

DEPARTMENT OF MATHEMATICS, PIEDMONT COLLEGE, 165 CENTRAL AVE., DEMOREST, GA 30535

E-mail address: mmontgomery@piedmont.edu